

AD-A045 508

SCHOOL OF AEROSPACE MEDICINE BROOKS AFB TEX
THERMAL RESPONSE IN 'MACACA MULATTA' EXPOSED TO 15- AND 20-MHZ --ETC(U)
SEP 77 J H KRUPP
SAM-TR-77-16

F/G 6/18


UNCLASSIFIED

NL

1 OF 1
ADA045508



END
DATE
FILMED
11-77
DDC




AD A045508

Report SAM-TR-77-16

K

12

**THERMAL RESPONSE IN *MACACA MULATTA*
EXPOSED TO 15- AND 20-MHZ RADIOFREQUENCY
RADIATION**



for C

September 1977

Interim Report for Period January-December 1976

Approved for public release; distribution unlimited.

AD No. _____
DDC FILE COPY

USAF SCHOOL OF AEROSPACE MEDICINE
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas 78235



NOTICES

This interim report was submitted by personnel of the Radiation Physics Branch, Radiation Sciences Division, USAF School of Aerospace Medicine, Aerospace Medical Division, AFSC, Brooks Air Force Base, Texas, under job order 7757-01-43.

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The animals involved in this study were maintained and used in accordance with the Animal Welfare Act of 1970 and the "Guide for the Care and Use of Laboratory Animals" prepared by the National Academy of Sciences-National Research Council.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

Jerome H. Krupp
 JEROME H. KRUPP, D.V.M.
 Project Scientist

John C. Mitchell
 JOHN C. MITCHELL, B.S.
 Supervisor

Robert G. McIver
 ROBERT G. MCIVER
 Brigadier General, USAF, MC
 Commander

| | |
|---------------------------------|---|
| ACCESSION for | |
| NTIS | White Section <input checked="" type="checkbox"/> |
| DDC | Blue Section <input type="checkbox"/> |
| UNANNOUNCED | <input type="checkbox"/> |
| JUSTIFICATION | <input type="checkbox"/> |
| BY | |
| DISTRIBUTION/AVAILABILITY CODES | |
| D. | SPECIAL |
| A | |

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|---|-----------------------|--|
| 1. REPORT NUMBER SAM-TR-77-16 | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) THERMAL RESPONSE IN MACACA MULATTA EXPOSED TO 15- AND 20-MHZ RADIOFREQUENCY RADIATION. | | 5. TYPE OF REPORT & PERIOD COVERED Interim Report, January-December 1976. |
| 7. AUTHOR(s) Jerome H. Krupp, D.V.M. | | 6. PERFORMING ORG. REPORT NUMBER |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS USAF School of Aerospace Medicine (RZP) Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas 78235 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62202F 7757-01-43 |
| 11. CONTROLLING OFFICE NAME AND ADDRESS USAF School of Aerospace Medicine (RZP) Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas 78235 | | 12. REPORT DATE September 1977 |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1213p. | | 13. NUMBER OF PAGES 10 |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | 15. SECURITY CLASS. (of this report) Unclassified |
| 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Radiofrequency Radiation Thermal Response | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Rhesus monkeys were exposed to high incident-power levels of 15- and 20-MHz radiofrequency radiation for 3 hours. After a modest rise in rectal temperature, thermoregulatory control was established. Temperature rise was related to frequency and incident power. Equivalent power absorptions in man at these frequencies would be 15-25 times greater than current personnel exposure limits. | | |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

THERMAL RESPONSE IN MACACA MULATTA EXPOSED TO 15- AND 20-MHZ RADIOFREQUENCY RADIATION

INTRODUCTION

Previous exposures of rhesus monkeys (Macaca mulatta) to 26-MHz radiofrequency (RF) radiation at incident power levels up to 1000 mW/cm² showed that the absorbed power produced a modest rectal temperature rise with compensatory thermoregulation which was maintained for the entire 6-hour exposure period (1). To further explore the high-frequency (HF) band, additional animals were similarly exposed to 15- and 20-MHz RF radiation at each of two different power levels.

METHODS AND MATERIALS

Ten rhesus monkeys, weighing 3.5 ± 0.5 kg, were housed in the vivarium, fed a commercial ration twice daily, and allowed water ad libitum. Each morning of the experiment, they were placed in slotted plastic cages, 20 x 20 x 40 cm, and transported to the RF-radiation facility. Following a short rest period, rectal and skin temperatures were taken, then two animals were placed in a rectangular coaxial HF-band transmission line (5). The remaining animals were kept adjacent to the exposure chamber, but outside the radiation field. Every half-hour rectal temperatures were recorded from each animal; the radiation was off for approximately 2 minutes at this time. The exposures continued for a total of 3 hours. Temperatures of all animals were recorded immediately after termination of exposure, and for the exposed pair, one additional recording was made 10 minutes post exposure. Two exposure periods were done each day until all animals had been exposed to one combination of frequency and power. This procedure was repeated until all animals had been exposed to four combinations, two frequencies at two power levels each.

Rectal temperatures were recorded using thermistor probes (Yellow Springs Instrument Telethermometer Model 73). Temperatures were estimated $\pm 0.1^\circ$ C. The thermistor was calibrated with an NBS-certified mercury thermometer (Taylor Instrument Co.).

All exposures were obtained in the HF-band TEM-mode exposure device, with two animals being exposed simultaneously. Power was supplied to the exposure device by an AN/FRT-6B transmitter operating in the continuous-wave (CW) mode and was maintained at the desired input power $\pm 4\%$. The 20-MHz exposures were made using input powers of 40 and 24 kW (CW), and the 15-MHz exposures were made with input powers of 40 and 29 kW. The E-fields were vertical, and animals were exposed in the erect position (E-polarization). Animals were exposed above the center conductor on the lateral center line: one animal 50 cm forward of and one 50 cm behind the longitudinal center line.

The E- and H-field maps of the cage were uniform within $\pm 2\%$. Measurements were obtained with calibrated 10-cm dipole and loop antennas read out on a Kiethley 600B electrometer.

Field dimensions are summarized in Table 1. Comparative values for man exposed to these conditions were obtained from the USAFSAM RF Radiation Dosimetry Handbook (3) and are shown in Table 2.

TABLE 1. DOSIMETRY DATA

| Freq. (MHz) | Input power (kW) | Average E-Field (V/m) | | Average H-Field (A/m) | | E/H ratio (ohms) | Incident power density ^a (mW/cm ²) |
|----------------|------------------------|--------------------------|--------------|--------------------------|--------------|------------------------|--|
| | | Front cage | Rear cage | Front cage | Rear cage | | |
| 20 | 40 | 2220 | 2160 | 4.85 | 4.75 | 450 | 1270 |
| | 24 | 1710 | 1680 | | | | 760 |
| 15 | 40 | 2020 | 1915 | | | 375 | 1025 |
| | 29 | 1750 | 1670 | | | | 775 |

^aCalculated from E-field data and averaged for cage position.

TABLE 2. COMPARATIVE EXPOSURE VALUES FOR MAN

| Primate Data | | For Equivalent Absorbed Power in Man | |
|------------------------|--|--|---|
| Exp. freq. (MHz) | Incident power (mW/cm ²) | Incident power at same freq. (mW/cm ²) | Freq. for same incident power (MHz) |
| 20 | 1270 | 225 | 8 |
| 20 | 760 | 150 | 8 |
| 15 | 1025 | 205 | 5 |
| 15 | 775 | 155 | 5 |

RESULTS AND DISCUSSION

Figures 1-4 show the mean rectal temperature differences for the four exposure conditions. Mean temperature differences and the standard errors are shown in Table 3. The differences were obtained by subtracting the mean of control temperatures for a given condition from the mean of exposed animals (Table 4). For each animal, control temperature means were derived from the control values that corresponded to his exposure time of day, thus eliminating the significant time-of-day effect which existed. Figure 5 shows the means of Table 4 for exposed animals through the full 190-minute observation period. Figure 6 gives the maximum temperature rise that occurred during the 3-hour period of exposure for the conditions of this experiment, together with data from previous work at 26 MHz.

The differences (exposed-control) show that all four irradiation conditions had a significant change in temperature over time ($P < .001$). In general, temperatures rose quickly in the first 30 minutes, then leveled off and fluctuated slightly or continued a slow rise for a time as thermoregulatory mechanisms were brought into play. By comparing all four conditions with regard to trend in temperature rise, some frequency X time interaction was indicated ($P < .08$). There is a suggestion that the higher frequency increases the body temperature more during the first hour and then maintains the higher level throughout the exposure period. This would conform to the concept of a thermal burden based on absorbed power (4), which has been shown to be frequency dependent (2). The homeostatic-temperature-control ability of the animals to adjust to the stress at levels 15 to 25 times greater than current permissible exposure levels for man, lends support to standards which are based in part on thermal burden and frequency. Following cessation of radiation, there was a significant drop in temperature within 10 minutes except for the 15 MHz/775 mW/cm² group ($P < .05$).

In summary, exposure of rhesus monkeys to RF radiation in the HF range, at power levels approximating 15-25 times the maximum permissible exposure limits for man, results in a modest rise in body core temperature which is adequately compensated for over an extended period of time, i.e., 3 hours or more. The rectal temperature response is related to frequency and field strength and conforms to generally accepted views regarding frequency, power absorption, and body size.




Figure 1. Mean rectal temperature differences for 20 MHz, 1270 mW/cm² exposure.




Figure 2. Mean rectal temperature differences for 20 MHz, 760 mW/cm² exposure.




Figure 3. Mean rectal temperature differences for 15 MHz, 1025 mW/cm² exposure.




Figure 4. Mean rectal temperature differences for 15 MHz, 775 mW/cm² exposure.

Table 3. MEAN TEMPERATURE ($^{\circ}\text{C}$) DIFFERENCES
(Irradiated - Control)

| Time (min) | 20/1270 ^a | 20/760 | 15/1025 | 15/775 |
|--------------------|----------------------|--------|---------|--------|
| 0 | .36 | -.30 | -.01 | -.15 |
| 30 | 1.13 | .42 | .81 | .29 |
| 60 | 1.38 | .56 | .65 | .34 |
| 90 | 1.30 | .92 | .85 | .41 |
| 120 | 1.57 | .76 | .88 | .71 |
| 150 | 1.60 | .88 | .75 | .79 |
| SE(D) ^b | .215 | .162 | .141 | .164 |

^aFrequency (MHz)/incident power (mW/cm^2).

^bStandard error of the mean difference. Not valid for comparing two differences.

TABLE 4. MEAN TEMPERATURES ($^{\circ}\text{C}$) OF EXPOSED PRIMATES

| Time (min) | 20/1270 ^a | 20/760 | 15/1025 | 15/775 |
|--------------------|----------------------|--------|---------|--------|
| 0 | 39.18 | 38.46 | 38.75 | 38.56 |
| 30 | 39.91 | 39.16 | 39.55 | 38.98 |
| 60 | 40.04 | 39.28 | 39.37 | 39.01 |
| 90 | 39.94 | 39.58 | 39.51 | 39.01 |
| 120 | 40.15 | 39.32 | 39.44 | 39.20 |
| 150 | 40.15 | 39.43 | 39.30 | 39.27 |
| 180 | 39.95 | 39.46 | 39.25 | 39.06 |
| Post 10 | 39.31 | 39.07 | 39.05 | 38.93 |
| SE(D) ^b | .203 | .144 | .083 | .129 |
| SE ^c | .170 | .104 | .036 | .065 |

^aFrequency (MHz)/incident power (mW/cm^2).

^bStandard error of the difference between two time means for the same radiation.

^cCalculated standard errors of the difference between 180 exposure minutes and 10 minutes post irradiation.




Figure 5. Mean rectal temperatures, all exposed animals.




Figure 6. Mean core temperature rise during three hours of exposures to 26, 20, and 15 MHz as a function of incident power.

REFERENCES

1. Frazer, J. W., et al. Thermal responses to high-frequency electromagnetic radiation fields. SAM-TR-76-20, Sep 1976.
2. Ghandi, P. P. Strong dependence of whole animal absorption and frequency of radio-frequency energy. Ann NY Acad Sci 247:532-578 (1975).
3. Johnson, C. C., et al. Radiofrequency radiation dosimetry handbook. SAM-TR-76-35, Sep 1976.
4. Michaelson, S. M., R. A. E. Thompson, and J. W. Howland. Biologic effects of microwave exposure. Rome Air Development Center TR-67-41, DDC No. AD 824-242, 1967.
5. Mitchell, J. C. A radiofrequency radiation exposure device. SAM-TR-70-43, July 1970.